



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Miss Knight remarked upon the mosses found in the dells of Wisconsin and announced the discovery by her of *Eustichium Norvegicum*, Br. Eu., in fruit.

Prof. Martin exhibited a pink variety of *Impatiens fulva*, Nutt., and an example of heterogamy in Indian corn.

At the regular meeting held Tuesday evening, Oct. 9th, the President occupied the chair and twenty persons were present.

Plants noted on Field Days.—Messrs. Wilber and Bicknell reported that they had found a white-rayed variety of *Aster Novæ-Angliæ*, L., at Van Cortlandt, N. Y. Mr. Hollick stated that he had found the var. *roseus* of the same species on Staten Island.

Dr. Britton gave Tottenville, S. I., as a station for *Trichostema lineare*, Nutt., a plant new to the Island.

Prof. Hyatt remarked upon some of the more notable plants that had been observed by him during a trip between Cincinnati and Chattanooga.

Adventitious Leaves in Rhus.—Mr. Schrenk exhibited a specimen of *Rhus Toxicodendron*, L., in which the inflorescence was replaced by tufts of leaves. (The same change has heretofore been noted in the BULLETIN as occurring in three other species of the genus—*R. glabra typhina* and *copallina*.)

At the meeting of Tuesday evening, Nov. 13th, the chair, in the absence of the presiding officers, was occupied by Prof. Day. There were twenty-six persons present.

Teratological.—Mr. Brower exhibited a specimen of *Amorpha fruticosa*, L., with fasciated stem, and one of *Cocculus Carolinus*, DC., with polymorphous leaves.

Mr. Schrenk remarked upon the structure of tuckahoe and upon the parasitism of *Gerardia tenuifolia*, Vahl (see page 132.)

Late-flowering Plants.—The following plants were reported by various members as being in flower for the second time this season: *Lonicera semipervirens*, Ait. (Miss Knight); *Viola sagittata*, Ait., *V. pedata*, L., and *V. primulæfolia*, L. (Mr. Hollick); *V. cucullata*, Ait. (Mr. Bicknell); and *V. lanceolata*, L. (Mr. Bisky). Dr. Britton remarked upon and exhibited under the microscope, specimens of *Protococcus (vulgaris?)* taken from the bark of trees in Brooklyn.

On motion the chairman appointed a committee of three (Messrs. Britton and Hyatt and Miss Knight) to consider a plan for forming a subsection of the Club for the study of physiological botany.

One person was elected an active member.

Correction.—*Pyrus arbutifolia* in Mr. Ridgway's note on pages 121 and 122 should of course read *P. Americana*. The error is so apparent that every reader of the BULLETIN has doubtless noticed and corrected it.